

Asymmetry of eye movement disturbances in schizophrenic patients

Alina Borkowska¹, Janusz K. Rybakowski²

¹Department of Psychiatry, University School of Medical Sciences, Bydgoszcz, Poland

²Department of Adult Psychiatry, University of Medical Sciences, Poznań, Poland

Thirty patients with paranoid schizophrenia were studied during exacerbation of symptoms. Control group consisted of thirty persons, age and sex-matched. Eye trackings were measured during point fixation and smooth pursuit movement. Frequency of intrusive saccades and total duration of square wave jerks were recorded for each eye. Schizophrenic patients had extensive eye tracking disturbances on both tasks, significantly more marked for the right eye than for the left one. In these patients, significant correlation was found between the intensity of the disturbance and the magnitude of right-left eye difference. The results point to the asymmetry of eye tracking disturbances in schizophrenia with greater right eye pathology.

Key words: schizophrenia, eye movements, lateralization, intrusive saccades, asymme-

Introduction

The eye movement disturbances in schizophrenia were first described in 1908 by Diefendorf and Dodge [11]. In 1970s [20] they were considered as a possible neurophysiological marker of vulnerability to schizophrenia. Such disturbances occur in 80% of patients with schizophrenia, in 50 % of their first degree relatives, and only in 10 % of healthy subjects [18]. While a genetic component is postulated [2, 19], the intensity of eye movement disturbances in schizophrenic patients may also be related to the acquired brain impairment [22, 24, 26].

The most frequent eye movement disturbances in schizophrenia include reduction of smooth pursuit gain, an increased number of intrusive saccades during smooth pursuit movement [19, 25, 33, 34], abnormalities of antisaccadic task, e.g. antisaccade latency and antisaccade errors rate [25, 29, 31, 39]. They also include abnormalities of visual fixation such as a decreased number of fixation points, a longer fixation time and an increased number of intrusive saccades occurring during fixation both in horizontal and vertical components [2, 35].

The data on the neuroanatomy of eye movements suggest that the areas of the brain involve mostly frontal and parietal cortical areas as well as the basal ganglia. Posterior

parietal areas specifically control visually guided eye movements, whereas frontal and parietal areas co-operate in controlling the more internally generated oculomotor functions [23]. Neuroimaging studies in schizophrenia showed that eye movement dysfunctions are connected with a decreased metabolism in the frontal and parietal association cortex [32]. In this study, patients with schizophrenia showing greater eye movement disturbances had significantly decreased metabolism in oculomotor region of frontal cortex. MacAvoy and Bruce [27] found an association between the intensity of eye movement disturbances and frontal impairment measured by neuroimaging and by the performance on Wisconsin Card Sorting Test.

According to the hypothesis of lateralization for cerebral dysfunctions, formulated by Flor-Henry [12], schizophrenic psychopathology is mostly associated with left hemisphere dysfunction. Recent results of neurophysiological and neuroimaging studies brought some confirmation of this concept. Studies with dichotic listening in schizophrenic patients point to abnormal right ear advantage, compatible with left hemisphere dysfunction [4, 21]. In paranoid schizophrenic patients, significantly more rightward conjugate lateral eye movement (CLEM) in response to emotional and spatial stimuli were described [38, 13, 3]. Neuroimaging studies in schizophrenia revealed structural changes of cortical and limbic regions, predominant on the left side [41, 5, 40, 10, 37] as well as disturbances of blood flow and glucose metabolism in left frontal and prefrontal cortex, especially during verbal tasks [14, 36]. Pathology of left cerebral hemisphere in schizophrenia has been accentuated during exacerbation of symptoms [16, 15, 4].

In this study, we put forward a hypothesis postulating that eye movement disturbances in schizophrenia during an acute episode (measured as intrusive saccades) may present an asymmetry, compatible with left hemisphere dysfunction, i.e. being more marked in the contralateral (right) eye. To this end, the eye movement disturbances measured in patients with schizophrenia were calculated separately for each eye and the data were compared with those of healthy control subjects.

Subjects

The experimental group consisted of thirty right-handed patients with paranoid schizophrenia (15 male and 15 female, aged 19-48 years, mean 31 ± 10 years). All subjects were hospitalised at an inpatient unit, Department of Psychiatry, University School of Medical Sciences in Bydgoszcz. None had the history of substance abuse, neurological and serious somatic illness or ophtalmic abnormalities. The diagnoses were made by psychiatric staff of the unit, using ICD-10 and DSM-IV criteria.

The assessments of eye movements were performed during exacerbation of psychotic symptoms, when patients were drug-free for at least 7 days. The mean intensity of schizophrenic symptoms as measured by PANSS scale at this time was 111 ± 20 points.

The control group consisted of thirty healthy right-handed persons, age and sex-matched (15 male and 15 female, aged 19-48 years, mean 33 ± 8 years). All subjects were somatically healthy and without any significant psychiatric history and without

any ophtalmic dysfunctions.

The project was accepted by the Ethical Committee at the University School of Medical Sciences in Bydgoszcz. All patients and control subjects gave the informed consent to the study, after the nature of the procedures had been fully explained to them.

Method

Eye trackings were measured by infrared reflectometry method, using Ober II system and Grenoble 96 computer program. Investigated subjects were seated in a darkened room, in front of the computer screen with goggles (with infrared detectors) on their eyes, the distance between 17-inches computer screen and subject's eye being 60 cm.

The conditions for measurements were sampling rate of 400 Hz and measurement time 20 sec, during two tasks: 1) point fixation on central position on the computer screen and 2) smooth pursuit at the moving point on Lisajou's curves. The reason for choosing the Lisajou curves in Ober II system for the measurement of smooth pursuit movement was that these curves have constant mathematical characteristics but, unlike sinusoidal curve, the subject investigated cannot predict the direction of the moving point.

The subjects were instructed to keep their eyes fixed on the point during point fixation task and to observe the moving point during smooth pursuit task.

The main indexes of eye movement disturbances were the frequency of intrusive saccades (IS) occurring during each task and total duration of square wave jerks (SWJ), regardless of their amplitude. The following assessment criteria were applied: 1) frequency of intrusive saccades (N/sec), 2) mean duration (SWJ) to spend eye away from the course of the observed point.

Blinks were detected and eliminated from the analysis. All results were calculated separately for the left eye and for the right one. For the frequency intrusive saccades and total duration of square wave jerks, the values of cut-off point (mean +1 SD for the results of healthy controls) were defined.

Statistical analysis was performed using Statistica 5.0 program. To evaluate normality distribution of the variables, the Shapiro-Wilk test was applied. Consequently, nonparametric tests were used for comparisons of differences between left and right eye (Wilcoxon test), between schizophrenic and control subjects (Mann-Whitney test) and between fractions of subjects (Chi-square test). Also Spearman correlation coefficients were calculated between the intensity of particular eye movement disturbances and the magnitude of right-left eye difference.

Results

The frequency of intrusive saccades as well as total duration of square wave jerks

Table 1

Frequency of intrusive saccades and total duration of square wave jerks (mean +SD) during point fixation and smooth pursuit calculated separately for left and right eye in patients with schizophrenia and in healthy controls.

| | | | LEFT EYE | RIGHT EYE |
|-------------------|---------------------------|---------------|------------|-------------|
| POINT FIXATION | Frequency of IS (/sec) | Schizophrenia | 0.45±0.25 | 0.95±0.5* |
| | | Controls | 0.14±0.15# | 0.16±0.14## |
| | Duration (msec) SJJ | Schizophrenia | 95±106 | 238±815** |
| | | Controls | 12±15## | 14±16## |
| SMOOTH PURSUIT | Frequency of IS (/sec) | Schizophrenia | 1.1±0.5 | 1.6±0.5* |
| | | Controls | 0.32±0.2# | 0.35±0.3# |
| | Duration (msec) SJJ | Schizophrenia | 358±813 | 634±1066** |
| | | Controls | 55±53## | 73±53## |

Difference right vs left eye * p<0.05 **p<0.01
Difference controls vs schizophrenia #p<0.01 ##p<0.001

during point fixation and smooth pursuit task, calculated separately for left and right eye in patients with schizophrenia and in healthy controls are shown in table 1.

Schizophrenic patients compared with healthy control subjects had significantly more eye tracking disturbances for all measures: intrusive saccades frequency and square wave jerks duration time, both in point fixation and smooth pursuit task, and for both left and right eye. In schizophrenic patients these disturbances were significantly more marked for the right eye than for the left one, for all measures on both point fixation and smooth pursuit task. In control subjects the values of disturbances were slightly higher for the right eye, however, never reaching significance vs. the left one.

A frequent finding in schizophrenic patients, especially during smooth pursuit task, was an appearance of intrusive saccade in the right eye, while at the same time the recording of the left eye was normal. This was never the case in control subjects. Such phenomenon was illustrated on figure 1.

In table 2, a percentage of schizophrenic patients with abnormal values (>cut-off point) during point fixation and smooth pursuit task in left and right eye is presented. The cut-off point was adopted as mean results + 1 SD for healthy control subjects.

In schizophrenic patients, the percentage of subjects with abnormal values (> cut-off point) for intrusive saccades frequency, and duration of square wave jerks amounted to 33-100 % on fixation task and to 47-100% on smooth pursuit task. These percentages were greater for the right eye on all measures. The difference between fractions of schizophrenic patients with abnormalities of brief intrusive saccades in right and left eye was significant, both during point fixation ($\chi^2=13.3$ p<0.001) and smooth pursuit task ($\chi^2=4.44$, p<0.0035). The most robust abnormalities, measured as values in schizophrenic patients falling outside cut-off point, were those occurring in the right eye. All patients with schizophrenia (100%) fell outside cut-off point for total duration of square wave jerks in the right eye, both during point fixation and smooth pursuit

Table 3

Correlation between the intensity of eye movement disturbances and the magnitude of right-left eye difference

| | | Spearman correlation coefficient | Significance |
|----------------|-------------------|----------------------------------|--------------|
| Point fixation | Duration SWJ msec | 0.636 | P=0.001 |
| | Frequency IS Msec | 0.522 | P=0.003 |
| Smooth pursuit | Duration SWJ msec | 0.524 | P=0.003 |
| | Frequency IS Msec | 0.274 | ns |

Correlation between the intensity of particular eye movement disturbances and the magnitude of right-left eye difference in schizophrenic patients is shown in table 3.

The correlation between the intensity of disturbances and right vs. left eye differences was highly significant for SWJ duration on both point fixation and smooth pursuit task, and for IS frequency on point fixation task.

Discussion

The results of this study confirm the existence of very profound abnormalities of eye tracking in schizophrenic patients during exacerbation of illness compared with control persons, in terms of highly increased frequency of intrusive saccades and duration of square wave jerks. This is in line with the results obtained by other researchers [25, 8, 31, 42]. However, the main finding of our study was the observation that these abnormalities in schizophrenic patients are clearly asymmetric, i.e. they are more marked in the right eye than in the left one. This may confirm our working hypothesis.

The studies on the possible laterality of eye movement disturbances in schizophrenia are scarce. Allen et al.[1] in their study of schizophrenic patients from Pacific population found a significantly higher number of antisaccadic errors for right visual field, pointing to dysfunction of the left hemisphere. Crawford et al. [9] found a correlation between a higher number of errors on antisaccadic task and a decreased blood flow in the anterior cingulate, insula and in the left striatum. Since the point fixation and smooth pursuit tasks used in this study did not involve right or left visual field, the asymmetry of eye movement disturbances found by us should predominantly be due to the differences in motor control of each eye exerted by the contralateral hemisphere.

In right-handed persons, the right eye performs the function of the "leading" eye and thus may be more vulnerable to the disturbances precipitated by external tasks. This was confirmed in our study by showing slight (statistically insignificant) asymmetry in eye movements observed in healthy persons (higher amplitude and higher

frequency of IS in right eye). However, in patients with schizophrenia, such asymmetry was highly significant and right eye movement disturbances in schizophrenic patients had the greater discriminant power versus healthy control subjects than such disturbances in the left eye. Our finding of a higher degree of right eye movement disturbances in schizophrenic patients may thus corroborate the results obtained in various studies, pointing to greater pathogenic involvement of the left hemisphere in schizophrenia [14, 40, 41].

High prevalence of eye movement disturbances in the first degree relatives of schizophrenic patients indicates a substantial genetic component to them. A locus on chromosome 6 has been postulated for these disturbances as a putative genotypic susceptibility marker to schizophrenia (Arolt et al., 1996). However, several lines of evidence point that the intensity of eye movement dysfunctions in schizophrenic patients, besides of genetic vulnerability, may be connected with acquired brain impairment. Kinney et al., [22] among subjects with eye movement disturbances found more common occurrence of obstetric complications in schizophrenic patients as compared to their first-degree relatives. In line with this, Malaspina et al. [26] found that the disturbances of smooth pursuit eye movement were more marked in non-familial schizophrenic patients than in patients with family history. Finally, Litman et al. [24] found a lower intensity of eye movement disturbances in monozygotic twin discordant for schizophrenia. To answer the question whether asymmetry of eye movement disturbances is connected with genetic or neurodevelopmental component one may speculate on the basis of our results concerning correlations between the intensity of particular eye movement disturbances and the magnitude of right-left eye asymmetry. Since highly significant correlations were obtained for all parameters measured on point fixation task and for total duration of square wave jerks on smooth pursuit task, it seems that the asymmetry of eye movement disturbances in schizophrenia may be mainly due to the brain impairment occurring in the course of neurodevelopmental process.

It would be of interest to know whether the asymmetries of eye movement disturbances found during exacerbation of symptoms in schizophrenia may persist into the period of clinical improvement and whether they also occur in 1st degree relatives of schizophrenic patients. Also, the possible effect of neuroleptic treatment on these dysfunctions may be worth studying. Finally, the research on this issue performed in other psychiatric illnesses (e.g. bipolar and unipolar mood disorders) where the opposite lateralization than in schizophrenia is reported [28, 6, 30, 3] may throw more light on this interesting phenomenon.

References

1. Alleen JS, Lambert AJ, Johnson FY, Schmidt K, Nero KL. *Antisaccadic eye movements and attentional asymmetry in schizophrenia in three Pacific populations*. Acta Psychiatr. Scand. 1996; 94: 258-265.
2. Arolt V, Lancer R, Nolte A, Muller-Myhsok B, Purmann S, Schurmann M, Leutelt J, Pinnow M, Schwinger E. *Eye tracking dysfunction is a putative phenotypic susceptibility marker of schizophrenia and maps to a locus on chromosome 6p in families with multiple occurrence of the disease*. Am. J. Med. Gen. 1996; 67: 564-579.
3. Borkowska A, Rybakowski J. *Conjugate lateral eye movements during exacerbation of paranoid*

- schizophrenia and the acute episode of major depression*. Biol. Psychiatry 1997; 41: 1065-1067.
4. Bruder G, Rabinowicz E, Towey J, Brown A, Kaufmann CA, Amador X, Malaspina D, Gorman J.M. *Smaller right ear (left hemisphere) advantage for dichotic fused words in patients with schizophrenia*. Am. J. Psychiatry 1995; 152: 932-935.
 5. Buschbaum MS, Yang S, Hazlett E, Siegel BV, Germans M, Haznedar M. *Ventricular volume and asymmetry in schizotypal personality disorder and schizophrenia assessed with magnetic resonance imaging*. Schizophr. Res. 1997; 27: 45-53.
 6. Cazard P, Ricard F, Facchetti L. *Depression and functional EEG asymmetry*. Ann. Med. Psychol. 1992; 150: 230-239.
 7. Clementz BA, McDowell JE. *Smooth pursuit in schizophrenia: Abnormalities of open-and closed-loop responses*. Psychophysiology 1994; 31: 79-86.
 8. Clementz BA. *Saccades to moving targets in schizophrenia: Evidence for normal posterior cortex functioning*. Psychophysiology 1996; 33: 650-654.
 9. Crawford TJ, Puri BK, Nijran KS, Jones B, Kennard C, Lewis SW. *Abnormal saccadic disractivity in patients with schizophrenia: a 99mTc-HMPO SPET study*. Psychol. Med. 1996; 26: 265-277.
 10. Deicken RF, Zhou L, Corwin F, Vinogradov S, Weiner MW. *Decreased left frontal lobe N-acetylaspartate in schizophrenia*. Am. J. Psychiatry 1997; 154: 688-690.
 11. Diefendorf AR, Dodge R. *An experimental study of the ocular reactions of the insane from photographic records*. Brain 1908; 31: 451-489.
 12. Flor-Henry P. *Psychosis and temporal lobe epilepsy: a controlled investigation*. Epilepsia 1969; 10: 363-395.
 13. Gur RE. *Left hemisphere dysfunction and left hemisphere overactivation in schizophrenia*. J. Abnorm. Psychology 1978; 87: 226-238.
 14. Gur RE, Jaggi JL, Shtasel DL, Ragland JD, Gur RC. *Cerebral blood flow in schizophrenia: effects of memory processing on regional activation*. Biol. Psychiatry 1994; 35:3-15.
 15. Green MF, Hugdahl K, Mitchel S. *Dichotic listening during auditory hallucinations in patients with schizophrenia*. Am. J. Psychiatry 1994; 151: 357-362.
 16. Harvey SA, Nelson E, Haller JW, Eary TS. *Lateralized attentional abnormality in schizophrenia is correlated with severity of symptoms*. Biol. Psychiatry. 1993; 33: 93-99.
 17. Holzman PS, Kringlen E, Levy DL, Proctor LR, Haberman S, Yasillo NJ. *Abnormal Pursuit eye movements in schizophrenia: evidence for a genetic indicator*. Arch. Gen. Psychiatry 1997; 34: 802-805.
 18. Holzman PS. *Recent studies of psychophysiology in schizophrenia*. Schizophr. Bull. 1987; 13: 65-91.
 19. Holzman PS, Solomon CM, Levin S, Wateraux CS. *Pursuit eye movement dysfunctions in schizophrenia: family evidence for specificity*. Arch. Gen. Psychiatry 1984; 41: 136-139.
 20. Holzman PS, Proctor LR, Hughes DW. *Eye tracking patterns in schizophrenia*. Science 1973; 181, 179-181.
 21. Karny N, Nachson I. *Abnormal lateralization in schizophrenia: empirical evidence for an integrated model*. Eur. Psychiatry 1995; 10: 75-84.
 22. Kinney DK, Levy DL, Yurgelun-Told D, Tramer S, Holzman P. *Inverse relationship of perinatal complications and eye tracking dysfunction in relatives of patients with schizophrenia: evidence for a two-factor model*. Am. J. Psychiatry 1998; 155: 976-978.
 23. Kömpf D, Heide W. *Cortical control of eye movements*. In: Kimura J, Shibasaki H, ed. *Recent Advances in Clinical Neurophysiology*. Amsterdam: Elsevier; 1996. p. 658-667.
 24. Litman RE, Torrey EF, Hommer DW, Radant AR, Pickar D, Weinberger DR. *A quantitative analysis of smooth pursuit eye tracking in monozygotic twins discordant for schizophrenia*. Arch. Gen. Psychiatry 1997; 54: 417-426.

25. Levy D, Holzman P, Matthyse S, Mendell N. *Eye tracking dysfunction and schizophrenia: a critical perspective*. Schizophr. Bull. 1993; 19: 461-536.
26. Malaspina D, Friedman JH, Kaufman C, Bruder G, Amador X, Strauss D. *Psychobiological heterogeneity of familial and sporadic schizophrenia*. Biol Psychiatry 1998; 43: 489-496.
27. MacAvoy MG, Bruce CJ. *Comparison of the smooth eye tracking disorder of schizophrenics with that of nonhuman primates with specific brain lesions*. Int. J. Neurosci. 1995; 80: 117-151.
28. Martinot JL, Hardy P, Feline A, Huret JD, Mazoyer B, Attar-Levy D, Pappata S, Syrota A. *Left prefrontal glucose hypometabolism in the depressed state*. Am. J. Psychiatry 1990; 147: 1313-1317.
29. McDowell JE, Clementz BA. *The effect of fixation condition manipulation on antisaccade performance in schizophrenia: Studies of diagnostic specificity*. Exp. Brain Res. 1997; 115: 333-344.
30. Min SK, Oh BH. *Hemispheric asymmetry in visual recognition of words and motor response in schizophrenic and depressive patients*. Biol. Psychiatry 1992; 31: 255-262.
31. Radant AD, Claypoole K, Wingerson DK, Cowley DS, Roy-Byrne PP. *Relationship between neuropsychological and oculomotor measures in schizophrenia patients and normal controls*. Biol Psychiatry 1997; 42: 497-805.
32. Ross D, Thaker G, Buchanan R, Kirkpatrick, Lahti A, Medoff D, Bartko JJ, Goodman J, Tien A. *Eye tracking disorder in schizophrenia is characterized by specific ocular motor defects and is associated with deficit syndrome*. Biol. Psychiatry 1997; 42: 781-796.
33. Rosse RB, Malthora AK, Kim SY, Deutsch SI. *Visual fixation deficits and evidence of cognitive impairment in schizophrenia*. Biol. Psychiatry 1992; 31: 412-416.
34. Rybakowski JK, Borkowska A. *Eye tracking abnormalities in schizophrenia during fixation and smooth pursuit movements*. Int. J. Psychophysiology 1998; 30: 171-172.
35. Rybakowski JK, Borkowska A, Araszkievicz A, Drózdź W, Kućma A. *The effect of risperidone compared with phenothiazine neuroleptics on eye tracking dysfunctions*. Psychiatria Danubina 1998; 10: 324.
36. Sachdev P, Brodaty H, Rose N, Haindl W. *Regional cerebral blood flow in late onset schizophrenia: a SPECT study using 99mTc-HMPAO*. Schizophr. Res. 1997; 27: 105-117.
37. Salisbury DF, Shenton ME, Sherwood AR, Fisher IA, Yurgelun-Todd DA, Tohen M. *First-episode schizophrenic psychosis and controls in P300 amplitude over left temporal lobe*. Arch Gen Psychiatry 1998; 55: 173-180.
38. Schweitzer L, Becker E, Welsh H. *Abnormalities of cerebral lateralization in schizophrenia patients*. Arch. Gen. Psychiatry 1978; 35: 982-985.
39. Sereno AB, Holzman PS. *Antisaccades and smooth pursuit eye movements in schizophrenia*. Biol. Psychiatry 1995; 37: 394-401.
40. Stefan MD, Murray RM. *Schizophrenia: developmental disturbance of brain and mind*. Acta Paediatr. 1997; Suppl. 422: 112-116.
41. Supprian T, Hoffman E, Warmuth-Metz M, Franzek E, Becker T. *MRI T2 relaxation times of brain regions in schizophrenic patients and control subjects*. Psychiatry Res. 1997; 75: 173-182.
42. Thaker GK, Ross DE, Cassady SL, Adami HM, LaPorte D, Medoff D, Lahti A. *Smooth pursuit eye movements to extraretinal motor signals. Deficits in relatives of patients with schizophrenia*.

Address for correspondence:

Alina Borkowska, Department of Psychiatry
University School of Medicine, Bydgoszcz
Kurpiskiego 19, 85-096 Bydgoszcz, Poland
e-mail: alab@aci.amb.bydgoszcz.pl

